

Boston Reed College
 Pharmacy Technician Program – A Comprehensive Approach
 Week #12 Lab

Refer to the labels on the following pages to fill in the information in the table below.

Brand	Generic	Dosage Strength	Form	Total Volume	Route	Manuft.	NDC #
	Furosemide						
	Dextrose and Sodium Chloride						
Kefzol							
	Lidocaine						
Nitrostat							
Vantin							
Robinul							
Oncovin							
Cleocin Phosphate							
Zithromax							
	Codeine						

In addition to the table above, put check mark (✓) next to the bar codes and circle the Control Numbers.

NDC 63323-280-02 28002

FUROSEMIDE

INJECTION, USP

20 mg/2 mL

(10 mg/mL)

For IM or IV Use Rx only

2 mL Single Dose Vial

Preservative Free

Discard unused portion.

PROTECT FROM LIGHT.

Do not use if discolored.

American Pharmaceutical Partners, Inc.

Los Angeles, CA 90024

401803A

LOT

EXP

1000 mL NDC 0074-7941-09

5% DEXTROSE and 0.9% SODIUM CHLORIDE Injection, USP

EACH 100 mL CONTAINS DEXTROSE, HYDROUS 5 g; SODIUM CHLORIDE 900 mg IN WATER FOR INJECTION. ELECTROLYTES PER 1000 mL: SODIUM 154 mEq; CHLORIDE 154 mEq, 560 mOsmo/LITER (CALC). pH 4.3 (3.5 to 6.5). ADDITIVES MAY BE INCOMPATIBLE. CONSULT WITH PHARMACIST, IF AVAILABLE. WHEN INTRODUCING ADDITIVES, USE ASEPTIC TECHNIQUE, MIX THOROUGHLY AND DO NOT STORE. SINGLE-DOSE CONTAINER. FOR I.V. USE. USUAL DOSAGE: SEE INSERT. STERILE, NONPYROGENIC. USE ONLY IF SOLUTION IS CLEAR AND CONTAINER IS UNDAMAGED. MUST NOT BE USED IN SERIES CONNECTIONS.

Rx only

3

LATEX

ABBOTT 2000 Ref. 59-0035-2/R4-10/99 PRINTED IN USA ABBOTT LABORATORIES, NORTH CHICAGO, IL 60064, USA

1
2
3
4
5
6
7
8
9

CAUTION—Federal (U.S.A.) law prohibits dispensing without prescription.

For I.M. or I.V. Use

Dosage—See literature.

To prepare solution add 2 mL Sterile Water for Injection or 0.9% Sodium Chloride Injection. Provides an approximate volume of 2.2 mL (225 mg per mL)

SHAKE WELL

Prior to Reconstitution: Store at Controlled Room Temperature 59° to 86°F (15° to 30°C)

Protect from Light

After Reconstitution: Store in a refrigerator. For Storage Time - See Accompanying Literature. If kept at room temperature, use within 24 hours.

Lyophilized

WV 4520 AMX

Eli Lilly & Co., Indianapolis, IN 46285, U.S.A.

Exp. Date/Control No.

NDC 0002-1497-01

VIAL No. 767

Rx

Lilly

KEFZOL®

STERILE CEFAZOLIN SODIUM, USP

Equiv. to

500 mg

Cefazolin

NDC 10019-017-56

Lidocaine HCl Injection, USP 1%

(10 mg/mL) Rx only

**FOR INFILTRATION AND NERVE BLOCK
NOT FOR SPINAL OR EPIDURAL ANESTHESIA**

30 mL Multiple Dose Vial

Baxter **ESILEDERLE™**

Mfd. for **Baxter Healthcare Corporation** affiliate
by: Elkins-Sinn, Cherry Hill, NJ 08003 400-741-01

Each mL contains lidocaine hydrochloride 10 mg, sodium chloride 7 mg and methylparaben 1 mg in Water for Injection. pH 5.0-7.0; sodium hydroxide and/or hydrochloric acid used, if needed, for pH adjustment.

Usual Dosage: See package insert for complete prescribing information.

Store at controlled room temperature 15°-30°C (59°-86°F).

Lot:
Exp.:



N 0071-0418-24

Nitrostat®

(Nitroglycerin Tablets, USP)

0.4 mg (1/150 gr)

Rx only

100 **SUBLINGUAL** TABLETS



PARKE-DAVIS

Warning—To prevent loss of potency, keep these tablets in the original container. Close tightly immediately after each use.

Usual Dosage—0.3 to 0.6 mg sublingually as needed.

See package insert for full prescribing information.

Keep this and all drugs out of the reach of children.

Dispense in original, unopened container.

Store at Controlled Room Temperature 20°-25°C (68°-77°F) [see USP].

Manufactured by:
Parke Davis Pharmaceuticals, Ltd.
Vega Baja, PR 00694

Distributed by:
PARKE-DAVIS
Div of Warner-Lambert Co
Morris Plains, NJ 07950 USA
© 1999, PDPL

6505-00-687-3663

0418G030

EXP:
LOT:

Rx only

See package insert for dosage and complete product information.

Warning: Not for injection

Store unconstituted product at controlled room temperature 20° to 25°C (68° to 77°F) [see USP]. Store constituted suspension in a refrigerator 2° to 8°C (36° to 46°F). Shake well before using. Keep container tightly closed. The mixture may be used for 14 days. Discard unused portion after 14 days.

Directions for mixing: Shake bottle to loosen granules. Add approximately 1/2 the total amount of distilled water required for constitution (total water = 29 mL). Shake vigorously to wet the granules. Add remaining water and shake vigorously.

Each 5 mL of suspension contains cefpodoxime proxetil equivalent to 100 mg cefpodoxime.

U.S. Patent Nos. 4,486,425; 4,409,215

Licensed from Sankyo Company, Ltd., Japan
Manufactured by
Pharmacia & Upjohn S.A.-N.V., Puurs - Belgium
for
Pharmacia & Upjohn Company
Kalamazoo, MI 49001, USA
817 150 101
5Q5358

NDC 0009-3615-03
50 mL (when mixed)

Vantin[®] For Oral Suspension

cefpodoxime proxetil
for oral suspension

100 mg per 5 mL

Equivalent to 100 mg per 5 mL
cefpodoxime when constituted



Pharmacia
& Upjohn

NDC 0031-7890-10

1 mL Single Dose Vial

ROBINUL[®]
INJECTABLE
(Glycopyrrolate
Injection, USP)

0.2 mg/mL

For I.M. or I.V. use

Pharmaceutical Division
A.H. Robins Company
Richmond, VA 23220

U7890-10-1

LOT

EXP.

NDC 0002-7194-01
1 mL VIAL No. 7194

Rx

Lilly

ONCOVIN[®]
SOLUTION
VINCRIStINE
SULFATE
INJECTION, USP

1

mg per
1 mL

FATAL IF GIVEN INTRATHECALLY
FOR INTRAVENOUS USE ONLY

Multiple Dose
REFRIGERATE

Eli Lilly, Indpls.

WW 1341 AMX

Exp. Date/Control No.

LOT/EXP

812728808

Single Dose Container
See package insert for complete product information. Store at controlled room temperature 20° to 25°C (68° to 77°F) (see USP). Do not refrigerate.
Pharmacia & Upjohn Co.
Kalamazoo, MI 49001, USA

NDC 0009-0870-21 2 mL Vial

Cleocin Phosphate®
clindamycin injection, USP

300 mg/2 mL

Equivalent to
300 mg clindamycin

Store at or below 86°F (30°C).

DOSAGE AND USE

See accompanying prescribing information.

Constitute to 100 mg/mL* with 4.8 mL of Sterile Water For Injection.

Must be further diluted before use.

For appropriate diluents and storage recommendations, refer to prescribing information.

*Each mL contains azithromycin dihydrate equivalent to 100 mg of azithromycin, 76.9 mg of citric acid, and sodium hydroxide for pH adjustment.

05-5191-32-0

CAUTION: Federal law prohibits dispensing without prescription.

NDC 0069-3150-83

Zithromax®
(azithromycin for injection)

For IV infusion only

STERILE
equivalent to

500 mg

of azithromycin
Distributed by



Pfizer Labs

Division of Pfizer Inc, NY, NY 10017

Lot No.

Exp. Date

See Package Insert for Complete Prescribing Information.

Store at Controlled Room Temperature 15°-30°C (59°-86°F).

PROTECT FROM MOISTURE.

Dispense in a well-closed container as defined in the USP/NF.

TABLETS IDENTIFIED 54 783

(Side One)  (Side Two) 

DO NOT USE UNLESS TABLETS CARRY THIS IDENTIFICATION

NDC 0054-4156-25

100 Tablets

EXP. LOT

30 mg 
CODEINE Sulfate Tablets USP

Each tablet contains Codeine Sulfate 30 mg
Rx only.

 **Roxane**
Laboratories, Inc.
Columbus, Ohio 43216

4151001
039
© RLI, 1999



3 0054-4156-25 7